

Table 8 Coral Reef Composition: Reef Extent

	DATA OPTION 1: Landsat ETM	DATA OPTION 2: MODIS
<i>Spatial Dimensions</i>		
Area to cover	185 km x 185 km per scene	Up to km ²
Example area	Heron Reef	
Mapping unit	15 m panchromatic 30 m multi-spectral	500 – 1000 m
Positional accuracy	Dependent on Geo-referencing process	Dependent on Geo-referencing process
<i>Temporal Dimensions</i>		
When	Approx 9.45 am	Approx 9.45 am
How often	every 16 days	Daily
Variable to map	Location and extent of reefs	Location and extent of reefs
Environmental / Sensor Restrictions	For sub-tidal vegetation to depth limited by water clarity. Inter-tidal and supra-tidal vegetation can have water on top. Not possible for turbid water Clouds, strong winds and breaking waves.	For sub-tidal vegetation to depth limited by water clarity. Inter-tidal and supra-tidal vegetation can have water on top. Not possible for turbid water Clouds, strong winds and breaking waves.
Processing technique (Output)	Supervised Image classification or feature detection, Object based analysis or manual delineation	Supervised Image classification
Resources – Hardware and Software	PC Image processing software GIS with image classification module (e.g. ARCGIS Image Analyst)	PC Image processing software with Hyper-spectral analysis capabilities, including sub- pixel mapping techniques.
Resource – Personnel	Trained in image classification Experience with Landsat data	Trained in image classification Experience with Landsat data

	Knowledge of reef geomorphology to be mapped When object based analysis then experience with object based analysis software (e.g. Ecognition)	Knowledge of reef geomorphology to be mapped hyper-spectral
References: Note these are some example references	(Ahmad et al., 1999; Andréfouët et al., 2005; Roelfsema and Phinn 2008; Phinn et al., in press)	(Andréfouët and Claereboudt 2000; Capolsini et al., 2003)

Ahmad, W., C. Menges and G. J. E. Hill (1999). Satellite Remote Sensing for monitoring reef zonation changes in Northern Australia. Proceedings of the 4th North Australian Remote Sensing and GIS Conference, Darwin, Australia.

Andréfouët, S. and M. Claereboudt (2000). "Objective class definitions using correlation of similarities between remotely sensed and environmental data." International Journal of Remote Sensing 21(9): 1925-1930.

Andréfouët, S., F. E. Muller-Karger, J. A. Robinson, C. J. Kranenburg, D. Torres-Pulliza, S. Spraggins and B. Murch (2005). Global assessment of modern coral reef extent and diversity for regional science and management applications: a view from space. 10th International Coral Reef Symposium, Okinawa, Japan, International Coral Reef Society, 1732-1745.

Capolsini, P., S. Andréfouët, C. Rion and C. Payri (2003). "A comparison of Landsat ETM+, SPOT HRV, Ikonos, ASTER, and airborne MASTER data for coral reef habitat mapping in South Pacific islands." Canadian Journal of Remote Sensing 29(2): 187–200.

Phinn, S. R., C. M. Roelfsema and P. J. Mumby (in press). "Multi-scale image segmentation for mapping coral reef geomorphic and benthic community zone." International Journal of Remote Sensing.

Roelfsema, C. M. and S. R. Phinn (2008). Evaluating Eight Field and Remote Sensing Approaches for Mapping the Benthos of Three Different Coral Reef Environments in Fiji. SPIE Asia-Pacific Remote Sensing, Noumea, SPIE, 14.